

Short communication

A novel de novo mutation in *LAMB3* causes localized hypoplastic enamel in the molar region

Kim Y-J, Shin TJ, Hyun H-K, Lee S-H, Lee ZH, Kim J-W. A novel de novo mutation in *LAMB3* causes localized hypoplastic enamel in the molar region. *Eur J Oral Sci* 2016; 124: 403–405. © 2016 Eur J Oral Sci

Amelogenesis imperfecta (AI) is a collection of diseases characterized by hereditary enamel defects and is heterogeneous in genetic etiology and clinical phenotype. In this study, we recruited a nuclear AI family with a proband having unique irregular hypoplastic pits and grooves in all surfaces of the deciduous molar teeth but not in the deciduous anterior teeth. Based on the candidate gene approach, we screened the laminin subunit beta 3 (*LAMB3*) gene and identified a novel de novo mutation in the proband. The mutation was a frameshift mutation caused by a heterozygous 7-bp deletion in the last exon (c.3452_3458delAGAAGCG, p.Glu1151Valfs*57). This study not only expands the mutational spectrum of the *LAMB3* gene causing isolated AI but also broadens the understanding of genotype–phenotype correlations.

Young-Jae Kim¹, Teo J. Shin¹, Hong-Keun Hyun¹, Sang-Hoon Lee¹, Zang H. Lee², Jung-Wook Kim^{1,3}

¹Department of Pediatric Dentistry & Dental Research Institute, School of Dentistry, Seoul National University, Seoul, Korea;

²Department of Cell and Developmental Biology & Dental Research Institute, School of Dentistry, Seoul National University, Seoul, Korea; ³Department of Molecular Genetics & Dental Research Institute, School of Dentistry, Seoul National University, Seoul, Korea

Dr Jung-Wook Kim, Department of Molecular Genetics, Department of Pediatric Dentistry & Dental Research Institute, School of Dentistry, Seoul National University, 275-1 Yongon-dong, Chongno-gu, Seoul 110-768, Korea

E-mail: pedoman@snu.ac.kr

Key words: enamel; genetic diseases; hereditary; *LAMB3*; mutation

Accepted for publication April 2016

Amelogenesis imperfecta (AI) is a collection of hereditary enamel defects that are heterogeneous in genetic etiology and clinical manifestation (1). Tooth enamel is the hardest calcified tissue in vertebrates, and its formation is under delicate control of a series of ectodermal–mesenchymal interactions (2). Any harmful influence, whether genetic or environmental, during amelogenesis could induce defects in the enamel. Frequently, an insult during development results in only enamel defects, indicating that amelogenesis is extremely vulnerable to such an event.

To date, more than 13 genes have been identified to cause AI. Even though the clinical phenotype is diverse and heterogeneous, some forms of AI share a common phenotype. Alternatively, there are some characteristic features related to a specific genetic etiology, enabling researchers to find the genetic cause based on the candidate gene approach (3). In the present study, we describe a family with AI with a unique pattern of irregular hypoplastic enamel pits and grooves. Mutational screening of the candidate gene successfully found a genetic etiology to associate genotype with phenotype.

Parents and a child with localized enamel hypoplasia visited the Department of Pediatric Dentistry, Seoul National University Dental Hospital. The parents and the child were informed of the study and agreed to participate. The study protocol was reviewed and approved by the Institutional Review Board of the Seoul

National University Dental Hospital. Clinical examinations and blood collections were performed after obtaining written consent, according to the Declaration of Helsinki.

The proband was a 6-yr-old boy from non-consanguineous parents. Both parents were healthy without any systemic and/or oral conditions. The pregnancy and delivery were uneventful. The boy had no past history of systemic illness or skin pathology. His deciduous dentition showed unique irregular hypoplastic enamel pits and grooves in all surfaces of his deciduous molars, but not in his anterior teeth (Fig. 1A). In particular, second molars exhibited altered crown shape as a result of severe enamel hypoplasia (Fig. 1B–E). However, the thin enamel showed a distinct contrast with the underlying dentin, indicating that mineralization was not severely affected. The patient's developing teeth also exhibited irregular enamel hypoplastic features (Fig. 1F).

The exons, including the exon–intron boundaries of the laminin subunit beta 3 (*LAMB3*) gene, were sequenced as previously described (4). Sanger sequencing of exon 23, the last exon of the *LAMB3* gene, revealed a heterozygous 7-bp deletion (c.3452_3458delAGAAGCG, p.Glu1151Valfs*57) (Figure S1). This deletion caused a shift in the reading frame, resulting in a change from glutamine to valine at codon position 1151 and the production of an additional novel 56 amino acids (thus giving 1207 amino




Fig. 1. Clinical photographs and panoramic radiograph of the proband. (A) Frontal clinical photograph of the proband. (B–E) Photographs showing occlusal and buccal surfaces of the deciduous molar teeth. All surfaces exhibit irregular hypoplastic enamel pits and grooves. The shape of the crown, especially that of the second deciduous molar, is altered as a result of severe enamel hypoplasia. (F) Panoramic radiograph of the proband. The irregular hypoplastic enamel shows a distinct contrast with the underlying dentin, indicating that enamel mineralization is not affected. The enamel of developing teeth also exhibits irregular hypoplasia.

acids in total), instead of the 1172-amino-acid wild-type LAMB3 protein. The parents were tested, but lacked the mutation, indicating that the mutation occurred spontaneously.

In this study, the proband exhibited enamel defects with unique irregular hypoplastic pits and grooves in all surfaces of primary molars but not in the anterior teeth. Hypoplastic AI can occur as a result of mutations in several genes: amelogenin, X-isoform (*AMELX*; MIM *300391) (5); enamelin (*ENAM*; MIM *606585) (6); family with sequence similarity 20 member A (*FAM20A*; MIM *611062) (7); integrin beta 6 (*ITGB6*; MIM *147558) (8); *LAMB3* (MIM *150310) (9–11); collagen type XVII alpha 1 (*COL17A1*; MIM *113811) (12); and laminin subunit alpha 3 (*LAMA3*; MIM *600805) (13).

Mutations in the *AMELX* gene are associated with X-linked hypoplastic and/or hypomaturational AI, and the clinical phenotype in male subjects is the generalized, rather than the localized, form of AI (1). Mutations in *ENAM* have been associated with generalized and localized hypoplastic enamel in an autosomal-dominant or autosomal-recessive manner. In the autosomal-recessive form, clinical severity has been shown to be related to the dosage of the mutant allele, and the carrier also exhibits some minor hypoplastic pits in several teeth (14). Frequently, horizontal hypoplastic grooves are a characteristic feature related to *ENAM*

mutations (15). Autosomal-recessive *FAM20A* mutations result in generalized hypoplastic AI with gingival hyperplasia, multiple eruption failures, and renal calcification (7, 16). Autosomal-recessive *ITGB6* mutations result in generalized hypoplastic or pitted hypomineralized AI (8, 17).

Based on the clinical appearance of the unique irregular hypoplastic pits and grooves of the proband, we selected the *LAMB3* gene for candidate gene screening. Mutational screening of the *LAMB3* gene successfully identified a novel mutation. If the candidate screening had revealed no pathologic variation in the *LAMB3* gene, the next targets would have been the *COL17A1* and *LAMA3* genes as a result of the similar clinical enamel phenotype, which can occur without skin problems even though this is extremely rare (12, 13, 18).

The genes *LAMB3*, *LAMA3*, and *COL17A1* are all involved in the skin disease, junctional epidermolysis bullosa (JEB) (19). Junctional epidermolysis bullosa is a rare inherited disorder featuring skin fragility and AI from defects in the genes encoding the components of hemidesmosome/basement-membrane complexes (20). Junctional epidermolysis bullosa is usually manifested as an autosomal-recessive inheritance pattern; however, in some cases, a heterozygous mutation can cause autosomal-dominant AI with little or no apparent skin defect (12, 13, 18). The finding of a novel mutation in this study will bring the number of *LAMB3* mutations causing isolated AI to seven (Table S1). All seven *LAMB3* mutations causing isolated AI are truncating ones that can be expressed by escaping the nonsense-mediated mRNA decay system. The observed enamel defects seem not to be caused by haploinsufficiency because most JEB carriers do not have any dental phenotype. Disease-causing *LAMB3* truncation mutations will act in a dominant-negative manner to cause enamel defects affecting developmentally vulnerable amelogenesis, without skin pathology.

Acknowledgments – We thank the participants in this study for their cooperation. This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korean government (2014R1A2A1A11049931).

References

1. WRIGHT T. Amelogenesis imperfecta. *Eur J Oral Sci* 2011; **119**(Suppl 1): 338–341.
2. HU JC, CHUN YH, AL HAZZAZZI T, SIMMER JP. Enamel formation and amelogenesis imperfecta. *Cells Tissues Organs* 2007; **186**: 78–85.
3. KANG HY, SEYMEN F, LEE SK, YILDIRIM M, TUNA EB, PATIR A, LEE KE, KIM JW. Candidate gene strategy reveals ENAM mutations. *J Dent Res* 2009; **88**: 266–269.
4. LEE KE, KO J, LE CG, SHIN TJ, HYUN HK, LEE SH, KIM JW. Novel LAMB3 mutations cause non-syndromic amelogenesis imperfecta with variable expressivity. *Clin Genet* 2015; **87**: 90–92.
5. LAGERSTROM M, DAHL N, NAKAHORI Y, NAKAGOME Y, BACKMAN B, LANDEGREN U, PETTERSSON U. A deletion in the amelogenin gene (AMG) causes X-linked amelogenesis imperfecta (AIH1). *Genomics* 1991; **10**: 971–975.
6. RAJPAR MH, HARLEY K, LAING C, DAVIES RM, DIXON MJ. Mutation of the gene encoding the enamel-specific protein,

- enamelin, causes autosomal-dominant amelogenesis imperfecta. *Hum Mol Genet* 2001; **10**: 1673–1677.
7. O'SULLIVAN J, BITU CC, DALY SB, URQUHART JE, BARRON MJ, BHASKAR SS, MARTELLI-JUNIOR H, DOS SANTOS NP, MAN-SILLA MA, MURRAY JC, COLETTA RD, BLACK GC, DIXON MJ. Whole-Exome sequencing identifies FAM20A mutations as a cause of amelogenesis imperfecta and gingival hyperplasia syndrome. *Am J Hum Genet* 2011; **88**: 616–620.
 8. POULTER JA, BROOKES SJ, SHORE RC, SMITH CE, ABI FARRAJ L, KIRKHAM J, INGLEHEARN CF, MIGHELL AJ. A missense mutation in ITGB6 causes pitted hypomineralized amelogenesis imperfecta. *Hum Mol Genet* 2014; **23**: 2189–2197.
 9. POULTER JA, EL-SAYED W, SHORE RC, KIRKHAM J, INGLEHEARN CF, MIGHELL AJ. Whole-exome sequencing, without prior linkage, identifies a mutation in LAMB3 as a cause of dominant hypoplastic amelogenesis imperfecta. *Eur J Hum Genet* 2014; **22**: 132–135.
 10. KIM JW, SEYMEN F, LEE KE, KO J, YILDIRIM M, TUNA EB, GENÇAY K, SHIN TJ, KYUN HK, SIMMER JP, HU JC. LAMB3 mutations causing autosomal-dominant amelogenesis imperfecta. *J Dent Res* 2013; **92**: 899–904.
 11. WANG X, ZHAO Y, YANG Y, QIN M. Novel ENAM and LAMB3 mutations in Chinese families with hypoplastic amelogenesis imperfecta. *PLoS ONE* 2015; **10**: e0116514.
 12. MCGARTH JA, CHRISTIANO AM, PULKINEN L, EADY RA, Uitto J. Compound heterozygosity for nonsense and missense mutations in the LAMB3 gene in nonlethal junctional epidermolysis bullosa. *J Invest Dermatol* 1996; **106**: 1157–1159.
 13. YUEN WY, PASMOOIJ AM, STELLINGSMA C, JONKMAN MF. Enamel defects in carriers of a novel LAMB3 mutation underlying epidermolysis bullosa. *Acta Derm Venereol* 2012; **92**: 695–696.
 14. ÖZDEMİR D, HART PS, FIRATLI E, AREN G, RYU OH, HART TC. Phenotype of ENAM mutations is dosage-dependent. *J Dent Res* 2005; **84**: 1036–1041.
 15. KIM JW, SEYMEN F, LIN BP, KIZILTAN B, GENÇAY K, SIMMER JP, HU JC. ENAM mutations in autosomal-dominant amelogenesis imperfecta. *J Dent Res* 2005; **84**: 278–282.
 16. WANG SK, AREF P, HU Y, MILKOVICH RN, SIMMER JP, EL-KHATEEB M, DAGGAG H, BAQAIN ZH, HU JC. FAM20A mutations can cause enamel-renal syndrome (ERS). *PLoS Genet* 2013; **9**: e1003302.
 17. WANG SK, CHOI M, RICHARDSON AS, REID BM, LIN BP, WANG SJ, KIM JW, SIMMER JP, HU JC. ITGB6 loss-of-function mutations cause autosomal recessive amelogenesis imperfecta. *Hum Mol Genet* 2014; **23**: 2157–2163.
 18. ALMAANI N, LIU L, DOPPING-HEPENSTAL PJ, LOVELL PA, LAICHEONG JE, GRAHAM RM, MELLERIO JE, McGRATH JA. Autosomal dominant junctional epidermolysis bullosa. *Br J Dermatol* 2009; **160**: 1094–1097.
 19. WRIGHT JT, FINE JD, JOHNSON L. Hereditary epidermolysis bullosa: oral manifestations and dental management. *Pediatr Dent* 1993; **15**: 242–248.
 20. MASUNAGA T. Epidermal basement membrane: its molecular organization and blistering disorders. *Connect Tissue Res* 2006; **47**: 55–66.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Table S1. Disease-causing mutations in the *LAMB3* gene.

Figure S1. Sequencing chromatograms of the proband and unaffected parents.